

SELECTED PUBLICATION LIST

1. Kasuga, M., Kahn, C.R., Hedo, J-A., Van Obberghen, E. and Yamada, K.M. Insulin-induced receptor loss in cultured human lymphocytes is due to accelerated receptor degradation. (1981) Proc. Natl. Acad. Sci. USA. 78: 6917-6921.
2. Kasuga, M., Karlsson, F.A. and Kahn, C.R. Insulin stimulates the phosphorylation of the 95K subunits of its own receptor. (1982) Science. 215: 185-187.
3. Kasuga, M., Van Obberghen, E., Nissley, P. and Rechler, M. Structure of the insulin-like growth factor receptor in chick embryo fibroblasts. (1982) Proc. Natl. Acad. Sci. USA. 79: 1864-1868.
4. Kasuga, M., Zick, Y., Blithe, D.L., Crettaz, M. and Kahn, C.R. Insulin stimulates tyrosine phosphorylation of the insulin receptor in a cell-free system. (1982) Nature. 298: 667-669.
5. Kasuga, M., Hedo, J-A., Yamada, K.M. and Kahn, C.R. The structure of insulin receptor and its subunits: Evidence for multiple nonreduced forms and a 210,000 possible proreceptor. (1982) J. Biol. Chem. 257: 10392-10399.
6. Kasuga, M., Fujita-Yamaguchi, Y., Blithe, D. and Kahn, C.R. Tyrosine-specific protein kinase activity is associated with the purified insulin receptor. (1983) Proc. Natl. Acad. Sci. USA. 80: 2137-2141.
7. Kasuga, M., Fujita-Yamaguchi, Y., Blithe, D.L., White, M.F. and Kahn, C.R. Characterization of the insulin receptor kinase purified from human placental membranes. (1983) J. Biol. Chem. 258: 10973-10980.
8. Kasuga, M., Sasaki, N., Kahn, C.R., Nissley, P. and Rechler, M.M. Antireceptor antibodies as probes of insulin-like growth factor receptor structure. (1983) J. Clin. Invest. 72: 1459-1469.

9. Kadowaki, T., Kasuga, M., Akanuma, Y., Ezaki, O. and Takaku, F. Decreased autophosphorylation of the insulin receptor-kinase in streptozotocin-diabetic rats. (1984) *J. Biol. Chem.* 259: 14208-14216.
10. Odawara, M., Kadowaki, T., Yamamoto, R., Shibasaki, Y., Tobe, K., Accili, D., Bevins, C., Mikami, Y., Matsuura, N., Akanuma, Y., Takaku, F., Taylor, S.I. and Kasuga, M. Human diabetes associated with a mutation in the tyrosine kinase domain of the insulin receptor. (1989) *Science* 245: 66-68.
11. Yamamoto, R., Koshio, O., Tobe, K., Shibasaki, Y., Momomura, K., Odawara, M., Kadowaki, T., Takaku, F., Akanuma, Y. and Kasuga, M. Phosphorylation state and biological function of a mutant human insulin receptor Val⁹⁹⁶. (1990) *J. Biol. Chem.* 265: 14777-14783.
12. Yonezawa, K., Yokono, K., Shii, K., Ogawa, W., Ando, A., Hara, K., Baba, S., Kaburagi, Y., Yamamoto-Honda, R., Momomura, K., Kadowaki, T. and Kasuga, M. In vitro association of phosphatidylinositol 3-kinase activity with the activated insulin receptor tyrosine kinase. (1992) *J. Biol. Chem.* 267:440-446.
13. Yonezawa, K., Ando, A., Kaburagi, Y., Yamamoto-Honda, R., Kitamura, T., Hara, K., Nakafuku, M., Okabayashi, Y., Kadowaki, T., Kaziro, Y. and Kasuga, M. Signal transduction pathways from insulin receptors to Ras: Analysis by mutant insulin receptors. (1994) *J. Biol. Chem.* 269: 4634-4640.
14. Kotani, K., Yonezawa, K., Hara, K., Ueda, H., Ido, Y., Sakaue, H., Ando, A., Chavanieu, A., Calas, B., Grigorescu, F., Nishiyama, M., Waterfield, M.D. and Kasuga, M. Involvement of phosphoinositide 3-kinase in insulin or IGF-1-induced membrane ruffling. (1994) *EMBO J.* 13: 2313-2321.
15. Ando, A., Yonezawa, K., Gout, I., Nakata, T., Ueda, H., Hara, K., Kitamura, Y., Noda, Y., Takenawa, T., Hirokawa, N., Waterfield, M.D. and Kasuga, M. A complex of GRB2-dynamin binds to tyrosine-phosphorylated insulin receptor substrate-1 after insulin treatment. (1994) *EMBO J.* 13: 3033-3038.
16. Hara, K., Yonezawa, K., Sakaue, H., Ando, A., Kotani, K., Kitamura, Y., Ueda, H., Stephens, L., Jackson, T.R., Hawkins, P.T., Dhand, R., Clark, A.E., Holman, G.D.,

- Waterfield, M.D. and Kasuga, M. Phosphoinositide 3-kinase activity is required for insulin-stimulated glucose transport but not for ras activation in CHO cells. (1994) Proc. Natl. Acad. Sci. USA. 91: 7415-7419.
17. Noguchi, T., Matozaki, T., Horita, K., Fujioka, Y. and Kasuga, M. Role of SH-PTP2, a protein-tyrosine phosphatase with Src homology-2 domains, in insulin-stimulated Ras activation. (1994) Mol. Cell. Biol. 14: 6674-6682.
18. Sakaue, M., Bowtell, D. and Kasuga, M. A dominant-negative mutant of mSOS1 inhibits insulin-induced ras activation and reveals ras-dependent and-independent insulin signaling pathways. (1995) Mol. Cell. Biol. 15: 379-388.
19. Fujioka, Y., Matozaki, T., Noguchi, T., Iwamatsu, A., Yamao, T., Takahashi, N., Tsuda, M., Takeda, T. and Kasuga, M. A novel membrane glycoprotein, SHPS-1, that binds the SH2 domain-containing protein tyrosine phosphatase SHP-2 in response to mitogens and cell adhesion. (1996) Mol. Cell. Biol. 16: 6887-6899.
20. Takada, T., Matozaki, T., Takeda, H., Fukunaga, K., Noguchi, T., Fujioka, Y., Okazaki, I., Tsuda, M., Yamao, T., Ochi, F. and Kasuga, M. Roles of the complex formation of SHPS-1 with SHP-2 in insulin-stimulated MAP kinase activation. (1998) J. Biol. Chem. 273: 9234-9242.
21. Tamori, Y., Kawanishi, M., Niki, T., Shinoda, H., Araki, S., Okazawa, H. and Kasuga, M. Inhibition of insulin-induced GLUT4 translocation by munc18c through interaction with syntaxin4 in 3T3-LI adipocytes. (1998) J. Biol. Chem. 273: 19740-19746.
22. Kotani, K., Ogawa, W., Matsumoto, M., Kitamura, T., Sakaue, H., Hino, Y., Miyake, K., Sano, W., Akimoto, K., Ohno, S. and Kasuga, M. Requirement of atypical PKC λ for insulin stimulation of glucose uptake but not for Akt activation in 3T3-LI adipocytes. (1998) Mol. Cell. Biol. 18: 6971-6982.
23. Takeda, H., Matozaki, T., Takada, T., Noguchi, T., Tuda, M., Ochi, F., Fukunaga, K., Inagaki, K. and Kasuga, M. PI 3-kinase γ and protein kinase C- ζ mediate RAS-independent activation of MAP kinase by a Gi protein-coupled receptor. (1999) EMBO J. 18: 386-395.

24. Noguchi, T., Matozaki, T., Inagaki, K., Tsuda, M., Fukunaga, K., Kitamura, Y., Kitamura, T., Shii, K., Yamanashi, Y., and Kasuga, M. Tyrosine phosphorylation of p62 Dok induced by cell adhesion and insulin: possible role in cell migration. (1999) *EMBO J.* 18: 1748-1760.
25. Takata, M., Ogawa, W., Kitamura, T., Hino, Y., Kuroda, S., Kotani, K., Klip, A., Gingras, A-C., Sonenberg, N., and Kasuga, M. Requirement for Akt (protein kinase B) in insulin-induced activation of glycogen synthase and phosphorylation of 4E-BP1. (1999) *J. Biol. Chem.* 274: 20611-20618.
26. Kitamura, T., Kitamura, Y., Kuroda, S., Hino, Y., Ando, M., Kotani, K., Konishi, H., Matsuzaki, H., Kikkawa, U., Ogawa, W., and Kasuga M. Insulin-induced phosphorylation and activation of cyclic nucleotide phosphodiesterase (PDE) 3B by the serine-threonine kinase Akt (1999) *Mol. Cell. Biol.* 19: 6286-6296.
27. Kawanishi, M., Tamori, Y., Okazawa, H., Araki, S., Shinoda, H., and Kasuga, M. Role of SNAP23 in insulin-induced translocation of GLUT4 in 3T3-L1 adipocytes. (2000) *J. Biol. Chem.* 275 : 8240-8247.
28. Inagaki, K., Yamao, T., Noguchi, T., Matozaki, T., Fukunaga, K., Takada, T., Hosooka, T., Akira, S., and Kasuga, M. SHPS-1 regulates integrin-mediated cytoskeletal reorganization and cell motility. (2000) *EMBO J.* 19:6721-6731.
29. Hosooka, T., Noguchi, T., Nagai, H., Horikawa, T., Matozaki, T., Ichihashi, M., Kasuga, M. Inhibition of the motility and growth of B16F10 mouse melanoma cells by dominant negative mutants of Dok-1. (2001) *Mol. Cell. Biol.* 21:5439-5449
30. Sakaue, H., Konishi, M., Ogawa, W., Asaki, T., Mori, T., Yamasaki, M., Takata, M., Ueno, H., Kato, S., Kasuga, M., Itoh, N. Requirement of fibroblast growth factor10 in development of white adipose tissue. (2002) *Genes Dev.* 16:908-912.
31. Matsumoto, M., Ogawa, W., Teshigawara, K., Inoue, H., Miyake, K., Sakaue, H., Kasuga, M. Role of the IRS-1 and PI 3-kinase signaling pathway in insulin-induced expression of SREBP-1c and glucokinase genes in rat hepatocytes. (2002) *Diabetes* 51:1672-1680.

32. Miyake, K., Ogawa, W., Mastumoto, M., Nakamura, T., Sakaue, H., Kasuga, M. Hyperinsulinemia, glucose intolerance, and dyslipidemia induced by acute inhibition of phosphoinositide 3-kinase signaling in the liver. (2002) *J. Clin. Invest.* 110:1483-1491.
33. Sakaue, H., Nishizawa, A., Ogawa, W., Teshigawara, K., Mori, T., Takashima, Y., Noda, T. and Kasuga, M. Requirement for 3-phosphoinositide-dependent kinase-1 (PDK-1) in insulin-induced glucose uptake in immortalized brown adipocytes. (2003) *J. Biol. Chem.* 278:38870-38874.
34. Matsumoto, M., Ogawa, W., Akimoto K., Inoue, H., Miyake, K., Furukawa, K., Hayashi, Y., Iguchi, H., Matsuki, Y., Hiramatsu, R., Shimano, H., Yamada, N., Ohno, S., Kasuga, M. and Noda, T. Protein kinase C λ in liver mediates insulin-induced SREBP-1c expression and determines both hepatic lipid content and overall insulin sensitivity. (2003) *J. Clin. Invest.* 112:935-944.
35. Inoue, H., Ogawa, W., Ozaki, M., Haga, S., Matsumoto, M., Furukawa, K., Hashimoto, N., Kido, Y., Mori, T., Sakaue, H., Teshigawara, K., Jin, S., Iguchi, H., Hiramatsu, R., Leroith, D., Takeda, K., Akira, S and Kasuga, M. Role of STAT-3 in regulation of hepatic gluconeogenic genes and of carbohydrate metabolism in vivo. (2004) *Nature Med.* 10:168-174.
36. Hashimoto N, Kido Y, Uchida T, Matsuda T, Suzuki K, Inoue H, Matsumoto M, Ogawa W, Maeda S, Fujihara H, Ueta Y, Uchiyama Y, Akimoto H, Ohno S, Noda T, and Kasuga M. PKC λ regulates glucose-induced insulin secretion through modulation of gene expression in pancreatic beta cells. (2005) *J. Clin. Invest.* 115:138-145.
37. Kanda H, Tamori Y, Shinoda H, Yoshikawa M, Sakaue M, Udagawa J, Otani H, Tashiro F, Miyazaki J, and Kasuga M. Adipocytes from Munc18c-null mice show increased sensitivity to insulin-stimulated GLUT4 externalization. (2005) *J. Clin. Invest.* 115:291-301.
38. Uchida T, Nakamura T, Hashimoto N, Matsuda T, Kotani K, Sakaue H, Hayashi Y, Nakayama KI, White MF, and Kasuga M. Deletion of *Cdkn1b* gene ameliorates hyperglycemia by maintaining compensatory hyperinsulinemia in diabetic mice. (2005) *Nature Med.* 11:175-182.

39. Hashimoto, N., Kido, Y., Uchida, T., Asahara, S., Shigeyama, Y., Matsuda, T., Takeda, A., Tsuchihashi, D., Nishizawa, A., Ogawa, W., Fujimoto, Y., Okamura, H., Arden, KC., Herrera, PL., Noda, T and Kasuga, M. Ablation of PDK1 in pancreatic β cells induces diabetes as a result of loss of β -cell mass. (2006) *Nature Genet.* 38:589-593.
40. Inoue, H., Ogawa, W., Asakawa¹, A., Okamoto, Y., Nishizawa, A., Matsumoto, M., Teshigawara, K., Matsuki, Y., Watanab², E., Hiramatsu, R., Notohara, K., Katayose, K., Okamura, H., Kahn, CR., Noda, T., Takeda, K., Akira, S., Inui, A and Kasuga, M. Role of hepatic STAT3 in brain insulin action on hepatic glucose production. (2006) *Cell Met.* 3:267-275.
41. Kanda, H., Tateya, S., Tamori, Y., Kotani, K., Hiasa, K., Kitazawa, R., Kitazawa, S., Miyachi, H., Maeda, S., Egashira, K and Kasuga, M. MCP-1 contributes to macrophage infiltration into adipose tissue, insulin resistance, and hepatic steatosis in obesity. (2006) *J. Clin. Invest.* 116:1494-1505.
42. Okamoto, Y., Ogawa, W., Nishizawa, A., Inoue, H., Teshigawara, K., Kinoshita, S., Matsuki, Y., Watanabe, E., Hiramatsu, R., Sakaue, H., Noda, T., Kasuga, M. Restoration of glucokinase expression in the liver normalizes postprandial glucose disposal in mice with hepatic deficiency of PDK1. (2007) *Diabetes.* 25 : 1000-1009.
43. Hosooka, T., Noguchi, T., Kotani, K., Nakamura, T., Sakaue, H., Inoue, H., Ogawa, W., Tobimatsu, K., Takazawa, K., Sakai, M., Matsuki, Y., Hiramatsu, R., Yasuda, T., Lazar, MA., Yamanashi, Y., and Kasuga, M. Dok1 mediates high-fat diet-induced adipocyte hypertrophy and obesity through modulation of PPAR- γ phosphorylation. (2008) *Nature Med.* 14:188-193.
44. Nishino, N., Tamori, Y., Tateya, S., Kawaguchi, T., Shibakusa, T., Mizunoya, W., Inoue, K., Kitazawa, R., Kitazawa, S., Matsuki, Y., Hiramatsu, R., Masubuchi, S., Omachi, A., Kimura, K., Saito, M., Amo, T., Ohta, S., Yamaguchi, T., Osumi, T., Cheng, J., Fujimoto, T., Nakao, H., Nakao, K., Aiba, A., Okamura, H., Fushiki, T., and Kasuga, M. FSP27 Contributes to efficient energy storage in murine white adipocytes by promoting the formation of unilocular lipid droplets.(2008) *J.Clin.Invest.*118:2808-2821.

45. Yasuda, K., Miyake, K., Horikawa, Y., Hara, K., Osawa, H., Furuta, H., Hirota, Y., Mori, H., Jonsson, A., Sato, Y., Yamagata, K., Hinokio, Y., Wang, H.Y., Tanahashi, T., Nakamura, N., Oka, Y., Iwasaki, N., Iwamoto, Y., Yamada, Y., Seino, Y., Maegawa, H., Kashiwagi, A., Takeda, J., Maeda, E., Shin, H.D., Cho, Y.M., Park, K.S., Lee, H.K., Ng, M., Ma, R., So, WY., Chan, J. Lyssenko, V., Tuomi, T., Nilsson, P., Groop, L., Kamatani, N., Sekine, A., Nakamura, Y., Yamamoto, K., Yoshida, T., Tokunaga, K., Itakura, M., Makino, H., Nanjo, K., Kadowaki, T. & Kasuga, M. Variants in KCNQ1 are associated with susceptibility to type 2 diabetes mellitus. (2008) *Nature Genet.*40:1092-1097.
46. Matsuda T, Kido Y, Asahara S, Kaisho T, Tanaka T, Hashimoto N, Shigeyama Y, Takeda A, Inoue T, Shibutani Y, Koyanagi M, Hosooka T, Matsumoto M, Inoue H, Uchida T, Koike M, Uchiyama Y, Akira S, Kasuga M. Ablation of C/EBP β alleviates ER stress and pancreatic β cell failure through induction of GRP78. (2010) *J. Clin. Invest.*120: 115-126.
47. Sakai M., Matsumoto M., Tujimural T., Yongheng C., Noguchi T., Inagaki K., Inoue H., Hosooka T., Takazawa K., Kido Y., Yasuda K., Hiramatsu R., Matsuki Y., Kasuga M. CITED2 links hormonal signaling to PGC-1 α acetylation in regulation of gluconeogenesis.(2012) *Nature Med.*18:612-617